



Emmanuel M.E. Billia
Júlia Zervanová

NEW *STEPHANORHINUS KIRCHBERGENSIS* (JÄGER, 1839) (MAMMALIA, RHINOCEROTIDAE) RECORDS IN EURASIA. ADDENDA TO A PREVIOUS WORK

NUOVE SEGNALAZIONI DI *STEPHANORHINUS KIRCHBERGENSIS*
(JÄGER, 1839) (MAMMALIA, RHINOCEROTIDAE) IN EURASIA.
ADDENDA AD UNA PUBBLICAZIONE PRECEDENTE

Riassunto breve - Viene proposto l'aggiornamento dei dati relativi alle segnalazioni di resti di *Stephanorhinus kirchbergensis* (JÄGER, 1839) in Europa e in Asia. La nota integra quanto già pubblicato (BILLIA 2011a) in precedenza.

Parole chiave: Rhinocerontidae, Pleistocene, *Stephanorhinus kirchbergensis*, Europa, Asia.

Abstract - An updated integration containing numerous other reports on *Stephanorhinus kirchbergensis* (JÄGER, 1839) remains in Europe and Asia is proposed here. The notes follow the previous paper (BILLIA 2011a).

Key words: Rhinocerontidae, Pleistocene, *Stephanorhinus kirchbergensis*, Europe, Asia.

Introduction

This work follows the previous one (BILLIA 2011a), recently published in *Acta Palaeontologica Romaniae*. It represents an updated integration containing several other data on sites (in both continents) which gave back *Stephanorhinus kirchbergensis* (JÄGER, 1839) remains. These data were acquired later based on further investigation (2011-2015).

Fig. 1 shows the present approximative distribution of the *S. kirchbergensis* Eurasian sites. Denmark apart (country not yet present in BILLIA 2011a), the paragraph numbers used here in the text correspond to the same used in the previous paper.

The present article is strictly correlated with the paper (BILLIA & ZERVANOVÁ in prep.) concerning the *S. kirchbergensis* coming from "Cava Italamenti" (Vernasso, Cividale del Friuli, Udine, Friuli Venezia Giulia, North-Eastern Italy) and other sites in adjacent areas where *S. kirchbergensis*, as well as other rhinoceros remnants, have been found throughout the time.

1. Europe

Denmark

A fragment of a left mandible (believed of Eemian age) from a gravel pit at Seest near Kolding (Southeast Jylland) would represent the only *S. kirchbergensis*

specimen ever found in Denmark (DEGERBØL 1952; AARIS-SØRENSEN 1988, 2009).

England, § 1.1, p. 19

The sequence at Crayford (Lower Thames Valley) (SCOTT 2009) shows great similarities to the Ilford one and has a similar research history (BRIDGLAND 1994; GIBBARD 1994). Just as at Ilford, the exact provenance of most specimens is unclear. The fauna from the Crayford Gravel underlying the brickearth includes *Stephanorhinus hemitoechus* (FALCONER) and *Coelodonta antiquitatis* (BLUMENBACH). The "lower brickearth" contained specimens of *S. kirchbergensis* and *C. antiquitatis* (SCHREVE 1997). Amino-acid ratios are indicative of a MIS 7 age for the temperate sediments at both sites (PENKMAN et al. 2008).

The Netherlands, § 1.2, p. 20

A *S. kirchbergensis* fragmentary second upper molar (M2), a calcaneum, and a third metacarpus (Mc III) - related with the Eemian interglacial - are reported for the first time from the Eurogeul (North Sea, off the coast of the Zuid-Holland province) (MOL et al. 2012).

Germany, § 1.4, pp. 20-22

Three other exceptionally well-preserved *Rhinoceros Merckii* (= *S. kirchbergensis*) teeth (a fourth upper premolar [29276], [Fig. 2, in this paper], a fourth lower premolar [G.S. 1795], and a second lower molar [G.S.

5120][Fig. 3, in this paper]) - described by HERMANN (1911: p. 14 - fig 1, p. 15 - figs 2, 3) - were found in the environs of Graudenz near Christburg (Westpreußen) (collections: at that time, Geologisch-Paläontologisch Institut der Universität Königsberg and Westpreußisch Provizial-Museum in Danzig).

Schroeder referred to some *Rhinoceros Merckii* (= *S. kirchbergensis*) remains from Jerxheim in Braunschweig (Niedersachsen) (SCHROEDER 1903) as well as from Heggen (Sauerland, Nordrhein-Westfalen) (SCHROEDER 1905).

Another *Rhinoceros Merckii* (= *S. kirchbergensis*) isolated tooth was found at Westeregeln (Magdeburg) (NEHRING 1878, 1880: p. 473; HERMANN 1911: p. 15).

A *Rhinoceros Merckii* (= *S. kirchbergensis*) second upper molar comes "aus den Sanden von Rixdorf bei Berlin" (POHLIG 1887: p. 801 - fig. 2).

S. kirchbergensis remains come also from the sand pits at Niederlehme ("Rixdorfer Horizont"), 3 km northeast of Königs Wusterhausen, north of the A10 Highway (N 52° 19' 30", E 13° 40' 45") (Landkreis Dahme-Spreewald, Brandenburg, about 40 km south of Berlin) (WINGS et al. 2009: pp. 71-74, figs 2). On the "Rixdorfer Horizont" *vide autem* in DIETRICH (1932), CEPEK (1986), HEINRICH (2002), HERMSDORF (2000, 2002), WINGS et al. (2009).

Dicerorhinus kirchbergensis (= *S. kirchbergensis*) remains also come from Husarenhof bei Besigheim

(Stuttgart, Baden-Württemberg) (STAESCHE 1941: Pl 5 - figs 4-6, Pl 6 and 7) and from Steinheim a. d. Murr (STAESCHE 1941: Pl 11). These last ones are also cited in the "Steinheim a. d. Murr Interglacial (Holstenian) faunal association" by CZARNETZKI (1983).

Two other deposits with *S. kirchbergensis* are cited in literature: Spardorf bei Erlangen (Nordbayern) where a distal portion of a lower tooth ascribed to *Rhinoceros cf. Merckii* Jäger (= *S. kirchbergensis*) was found (Krumbeck 1917) and Schöningen (Ost-Niedersachsen) where both *S. kirchbergensis* and *S. hemitoechus* remains were recovered (BERKHOLST 2011: pp. 39-45, figs 14-16, tabs 3-4).

From "einer Höhle bei Heggen (Westfalen)" come *Rhinoceros Merckii* (= *S. kirchbergensis*) remains: a maxilla with P3-M2, a first upper premolar and some other odontological remains (REEKER 1907: p. 31) (conservation, at least at that time: Museum des Vereins für Heimatkunde des Süderlandes, Altena).

In BILLIA (2011a), the author incidentally omitted of reporting the WÜST (1901) paper, in which the *S. kirchbergensis* remains from Rabutz bei Gröbers, from the "Kalktuff von Weimar", from the "Kalktuff von Taubach bei Weimar" as well as the "*Rhinoceros?Merckii* Jäg." (= *S. kirchbergensis*) remains from Bilzingsleben bei Kindelbrück are widely described as well as figured (WÜST 1901: pp. 265-281, Pls IV and V).

According to LANSER (1997) "Bei Grabungsarbeiten in einem östlichen Seitengang der Dechenhöhle bei



Fig. 1 - A map showing the *S. kirchbergensis* Eurasian approximative distribution (also available on internet: http://commons.wikimedia.org/wiki/File:Distribution_of_S._kirchbergensis2.png) drawn after: BILLIA 2008, 2011a; BILLIA & PETRONIO 2009 and VAN DER MADE 2010.

- Mappa con la distribuzione euroasiatica indicativa di *S. kirchbergensis* (disponibile anche su: http://commons.wikimedia.org/wiki/File:Distribution_of_S._kirchbergensis2.png) disegnata sulla base di BILLIA 2008, 2011a; BILLIA & PETRONIO 2009 e VAN DER MADE 2010.




Fig. 2 - *S. kirchbergensis* from Graudenz, Germany: fourth upper premolar (after HERMANN 1911).
- *S. kirchbergensis* proveniente da Graudenz, Germania: quarto premolare superiore (da HERMANN 1911).




Fig. 3 - *S. kirchbergensis* from Graudenz, Germany: fourth lower premolar and second lower molar (after HERMANN 1911).
- *S. kirchbergensis* proveniente da Graudenz, Germania: quarto premolare inferiore e secondo molare inferiore (da HERMANN 1911).

Iserlohn-Letmathe (nördliche Sauerland), wurde im Frühjahr 1993 ein Rhinoceros schädel entdeckt. Die Grabungen wurden außerhalb des Schauhöhlenbereichs durchgeführt, der seinerzeit als Bodendenkmal eingetragen war" (Fig. 4, in this paper).

Italy, § 1.7, p. 23

According to STEHLIN (1930), the rhinoceros remains from the Leffe basin (Bergamo) belong to *Stephanorhinus*



Fig. 4 - Skull of *S. kirchbergensis* from the Dechenhöhle near Iserlohn-Letmathe (Germany), after LANSER (1997).
- Cranio di *S. kirchbergensis* rinvenuto nella Dechenhöhle presso Iserlohn-Letmathe (Germania), da LANSER (1997).

etruscus (Falconer). On the Leffe basin, *vide autem* in TARAMELLI (1898), VENZO (1953-56), VIALLI (1956), LONA (1963a, 1963b), BREDA & MARCHETTI (2003, 2007), RAVAZZI et al. (2009).

In July 2013 (PERSICO et al. 2014), an exceptionally well-preserved *S. kirchbergensis* skull (only the left third upper molar is missing) was discovered on alluvial sediments along the Po near Spinadesco (Cremona, Lombardia, Northern Italy). The skull does not exhibit abrasive signs, conditions which are in accordance with a short fluvial transport. Presently, the specimen is preserved in the "Museo Paleoantropologico del Po" at San Daniele Po (about 12 Km south-east of Cremona, Cremona province).

Slovenia, § 1.8, p. 24

The *D. kirchbergensis* (= *S. kirchbergensis*) remains from Kamnitnik (Skofija Loka, about 20 km northwest of Ljubljana) reported by RAKOVEC (1942: p. 247-250, Pl I - fig 5) consist of a second upper molar.

Croatia, § 1.9, p. 24

Mistakenly, the author (BILLIA 2011a) previously placed Volosko on the Lussino (Losinj) island (*vide p. 24*). Actually, Volosko and the Lussino island represent two different localities. Volosko/Volosco/Volosca (or Preluk/Preluka, such as in FABIANI 1919) lies north of Opatija/Abbazia (Kvarnar/Quarnaro gulf, Istria, Croatia). According to FABIANI (1919: p. 90), from this locality comes a first upper molar (FABIANI 1919: Pl XVII - figs 8a, 8b; Valle private collection, Trieste) ascribed by the same author to *Rhinoceros mercki* (= *S. kirchbergensis*). Later, MALEZ (1961: p. 77, 1963: pp. 112, 179) *sic et sempliciter* mentioned this toponym only. Afterwards, in another paper regarding a systematic revision of the whole Yugoslavian fauna, Volosko is not taken into consideration.




Fig. 5 - *Dicerorhinus kirchbergensis* (JÄGER, 1839) from Bizek (Medvednica Mountains, Zagreb, Croatia), second upper molar, ca. 0,9 natural size; 1: norma occlusalis, 2: norma pulparis (after MALEZ 1961).
 - *Secondo molare superiore di Dicerorhinus kirchbergensis* (JÄGER, 1839) da Bizek (Monti Medvednica, Zagabria, Croazia), ca. 0,9 grandezza naturale; 1: norma occlusalis, 2: norma pulparis (da MALEZ 1961).

On the Lussino/Losinj island (Kvarnar/Quarnaro archipelago, Dalmatia, Croatia) a second and a third upper molars (MCSN-TS s.n.) were found - in excellent conditions - in the first half of the XX century (LEONARDI 1945-47: p. 150-151, Pl VI - fig 1) (collections: Museo Civico di Storia Naturale, Trieste).

An exceptionally well-preserved lower molar (possibly a third) of unknown origin (LEONARDI 1945-47: pp. 151-152, Pl V - figs 1, 3) is showing a fossilization status very close to the two above mentioned molars.

Furthermore, because of both its morphological and biometrical features might belong to the same individual, so that we may assume that it comes from Lussino as well (LEONARDI 1945-47: p. 151).

As to the *S. kirchbergensis* material found in this area (Istria, Kvarnar/Quarnaro, Dalmatia) an overview is available in BILLIA (2011b).

The entire stock of rhinoceros remains collected at Krapina (GORJANOVIĆ-KRAMBERGER 1913b) - belonging to both *S. kirchbergensis* and *C. antiquitatis* - consists of 324 specimens in total.

The *Dicerorhinus mercki* (= *S. kirchbergensis*) third phalanx of Riss-Würm age from the Veternica cave (southwest of Medvednica, Medvednica Mountains, Zagreb, 306 m asl, 45° 50' 36" - 13° 32' 24") is described as well as figured in MALEZ (1958: pp. 5, 19, Pl I - figs 2a, 2b as well as in MALEZ 1963b: p. 109, Pl. XXII). This cave is situated ca 1,5 km north of Bizek. On the cave stratigraphy/speleology/geo-interpretations, vide autem in MALEZ (1963a).

MALEZ (1961: p. 66) also refers to a *D. kirchbergensis* second upper molar of large dimensions belonging to a young-adult individual found at Bizek (Medvednica Mountains), ca 1,5 km south of Veternica and about 9 km west of Zagreb. The exceptionally well-preserved tooth (s.n.) (Pl 1 - figs 1, 2; Pl 2 - figs 1, 2, 3; Pl 3 - fig 1) Fig. 5, in this paper) was found in a calcareous rock crack. MALEZ (1961: p. 65) correlated this upper molar with the Riss-Würm Interglacial (just as for Veternica, MALEZ 1961: p. 65). The remains are preserved at the Palaeontological Museum of JAZU in Zagreb.

Some other remains ascribed to this taxon could come from the Betalov Spodmol cave near Postojna (RAKOVEC 1959) (Riss-Würm Interglacial) and from the ossiferous breccie near Pula (MALEZ 1963b).

The four rhinoceros remains from Veternica (a left M2 [VTR. 199], a first phalanx [VTR. 200], a third phalanx [VTR. 201], and a rib shaft fragment [VTR. 202]), coming from the "j" level, were previously described by MALEZ (1963a: p. 108-112, Pl XXII, Pl XXIV - figs 1a-c) who assigned them to *D. kirchbergensis* (= *S. kirchbergensis*). Later, MIRACLE & BRAJKOVIC (1992: p. 2) "suggest that a more prudent identification of these remains is *Dicerorhinus* sp."

As to the rhinoceros remains from Hvar (WOLDRICH 1882: pp. 455-456; LEONARDI 1947: pp. 17-18), according to TOULA (1902: p. 5 and 1907: pp. 447-448) must be ascribed to *Rhinoceros etruscus* (= *S. etruscus*), not to *S. kirchbergensis*.

From the Vindija cave near Donja Voca (a cave in Tortonian conglomerated limestone, localised ca 55 km northeast of Zagreb and ca 20 km west of Varazdin) possibly come other *Dicerorhinus mercki* (= *S. kirchbergensis*) remains (VUKOVIĆ 1954: pp. 27-28). The cave was described in a previous work by the same author (VUKOVIĆ 1953).




Fig. 38. Vollkommen entwickelter, halb abgenützter Backenzahn von *Atelodus Merckii*, aus Rheinpreussen (vom $\frac{1}{14}$ verkleinert). el Aussenfalte (*Ectoloph*), ml Hinterfalte (*Metaloph*), pl Vorderfalte (*Protoloph*), ca Sporn (*Crista*), welchen hier *Parastelidion* vertritt, et *Antistetidion* (*Anticrochet*), Stelidion oder Crochet auf der Vorderfalte ist nicht entwickelt, pj vordere Grube, sj Mittelgrube, zh hintere Grube, g Cingulum, zh Hinterhöcker, ph Vorderhöcker, mit welchem die Aussenfalte z., wo die Vorderfalte endet.

Fig. 6 - Second upper molar of *Atelodus Merckii* (ancient synonym of *S. kirchbergensis*) from "Rheinpreussen" (after KAFKA 1913).
- Il secondo molare superiore di *Atelodus Merckii* (sinonimo di *S. kirchbergensis*) da "Rheinpreussen" (da KAFKA 1913).

Vaternica (MALEZ 1958: p. 7; SMITH 1976a; WOLPOFF 1979) and Vindija (MALEZ et al. 1980; WOLPOFF 1980; WOLPOFF et al. 1981) - just as for Krapina (GORJANOVIĆ-KRAMBERGER 1904, 1906, 1913a) and Velika Pećina (Ravna Gora) (SMITH 1976a, 1976b) - yielded significant well-known Late Pleistocene fossil hominids (JELINEK 1969; WOLPOFF et al. 1981, *inter alios*) as well as a lot of Late Mousterian-associated artefacts (VALOCH 1968, *inter alios*). Patterns of faunal/climatic changes were provided by MALEZ (1978a, 1978b, 1978c). By the way, another cave with significant Palaeolithic human traces is Crvena Stijena (Crna Gora/Montenegro) (RAKOVEC 1958). *Homo neandethalensis* King, 1864 remains were collected in Krapina, while Vaternica and Velika Pećina yielded *Homo sapiens* L., 1758 remains.

Czech Republic, § 1.11, p. 24

MOSTECKY (1966) reported on some *D. kirchbergensis* (= *S. kirchbergensis*) remains from the "Chlupáčova Služ"

Cave near Koněprusy (Beroun, southwest of Prague, Central Bohemia), a part of the largest Bohemian cave system - Koněpruské jeskyně (about 2 km in length and about 70 m vertical range), in a limestone region known as "Bohemian Karst". Some *C. antiquitatis* remains were also found into the same cave. However, the *D. kirchbergensis* remnants seem actually to belong to *S. hemitoechus*. As far as to concern an overview on Bohemian Karst, *vide autem* in DIEDRICH & ZAK (2006).

In KAFKA (1913: fig. 38), the *Atelodus Merckii* (ancient synonym of *S. kirchbergensis*) second upper molar from "Rheinpreussen" is figured (Fig. 6, in this paper).

According to some authors (TYRÁČEK et al. 2001, 2004) from Račiněves (Litoměřice okres, Ústecký kraj [Ústi region], Northern Bohemia) would come some *S. kirchbergensis* remains. Actually, *S. kirchbergensis* is listed in a faunal table only.

Slovakia, § 1.12, p. 24

From fluvial deposits along the Váh river at Šal'a (about 60 Km east of Bratislava, Šal'a district, Nitra region, Southwestern Slovakia) comes a right branch of a mandible with five very well-preserved teeth (P2-M2) (SCHMIDT 1969; ĎURIŠOVÁ 1994).

Poland, § 1.13, pp. 24-25

On p. 25 of BILLIA (2011a), GÜRICH (1908) is erroneously cited as a reference. Actually, on page 52 "... ein linker Unterkieferast von *Rhinoceros* sp. (von der Symphise an 5 Zähne umfassend) gefunden worden..." is the only rhinoceros remains cited in all the paper. However, the publication year is 1905 (non 1908) and *S. kirchbergensis* remains are completely absent. The five *Rhinoceros Merckii* Jäger (= *S. kirchbergensis*) teeth collected by Slósarski between 1880 and 1882 at Szczeslewice near Warsaw (SLÓSARSKI 1884: pp. 357, 370; Pls XIII and XIV) consist of a second upper molar and a first upper molar (Pl XIII - figs 6, 7), a fourth upper premolar, a second and a first upper molars (XIV - figs 5, 8, 9)(three reproductions of them [Pl XIV - fig 5 and Pl XIII - figs 5, 6] from SLÓSARSKI [1884] are in Fig. 7 of this paper).

Moldova, § 1.15, p. 25

The right hemimandible (OPS 1640) is also described in BELYAEVA & DAVID (1975: pp. 164-167, fig 4 and Pl II - fig 2). The first who reported the remains was SINZOW in 1900.

Ukraine, § 1.16, pp. 25-26

From a terrace on the left bank of the Kuyal'nik and Khadzhibeev estuaries (north of Odessa, Ukraine) together with *Archidiskodon wüsti* Pavlova, *Bison schoetensacki* Freudenberg, *Bison* aff. *priscus* Bojanus remains outcropped some *S. kirchbergensis* remnants




Fig. 7 - *Rhinoceros Merckii* JÄGER from Szczeslewice near Warsaw (Poland). 5 - fourth upper premolar, 6 - second upper molar, 7 - first upper molar (after SLÓSARSKI 1884).
- *Rhinoceros Merckii* JÄGER da Szczeslewice presso Warsaw (Polonia). 5 - quarto premolare superiore, 6 - secondo molare superiore, 7 - primo molare superiore (da SLÓSARSKI 1884).

as well (STEPANOV 1961; cited also in CHEPALYGA 1967: p. 48).

Russian Federation (European area), § 1.17, p. 26

SATAEV (2008: p. 144) refers to a *Stephanorhinus* sp. third deciduous molar crown (*S. kirchbergensis*, according to the author) from the "Verkhnyaya" Cave (Southern Ural).

In 2012, on a bank of the lower course of the Volga at Cherny Yar (Astrakhan' oblast') a fourth metacarpal (Mc IV) probably belonging to *S. kirchbergensis* has been recovered (unpublished material, pers. communication). From Cherny Yar (in literature also known as "Nizhnee Zaymishche") come other well-known *S. kirchbergensis* material: the full mandible [ZIN 16948] and the right hemimandible [ZIN 29854] described by GROMOVA (1935: pp. 98-101) (*vide autem* in BILLIA 2008: p. 28, fig 2-b, c; 2014: p. 186).

For a new, detailed description concerning all the other Russian-european sites which gave back *S. kirchbergensis* remains *vide autem* in BILLIA (2014).

2. Asia

Russian Federation (Asian area), § 2.1, p. 26

FRIEDEL (1880: pp. 353-354) as well as SCHRENCK (1880: pp. 1-2) referred to a find of *R. Merckii* Jäger (= *S. kirchbergensis*) remains on a bank of the Butantai river (a tributary of the Jana river) in the Vekhoyansky district (North-Eastern Siberia) occurred in 1877. According to CHERSKY (1878: p. 165, 1879: pp. 36-59), those remains must be assigned to *C. antiquitatis*.

The two teeth from Yakutya, previously described by DUBROVO (1957), were later also reported by ALEKSEEV (1970).

In 2013, along the Chumysh river (Kytmanov district, Altaysky kray, Central-southern Siberia) five *S. kirchbergensis* teeth (two fourth upper premolars, two upper molars and a third upper molar) together with a juvenilis mandibular left branch with three deciduous molar (dp2-dp4), and a mandibular right branch with five teeth (P2-M3) have been found. Moreover, along the Chik river (Kochenev district, Novosibirsk region, Central-southern Siberia) a *S. kirchbergensis* third upper molar has also been recovered (LYUBACHEV et al. 2014).

Furthermore, other *S. kirchbergensis* teeth - of remarkable dimensions - come from three other Central-Siberian localities situated along the Ob' and the Irtysh (unpublished material, pers. communication). Throughout the time, other *S. kirchbergensis* material outcropped along both these rivers (ALEKSSEVA 1980; SHPANSKY & BILLIA 2012, *inter alios*).

Azerbaijan, § 2.3, p. 26

More on the Azykh cave. The cave (about 900 m asl) - a six-cave complex known as a living site of stone-age man - lies about 3 km northeast of the Tugh village in Nagorno-Karabakh (at the boundary with South Azerbaijan where the Araks river cuts the slopes of the Lesser Caucasus and the Mil-Mugan steppes) on the left bank of the Kuruchay river. It was found by M. Guseinov in 1960. Among some others, systematic investigation have been carried out there by A.V. Mamedov and S.D. Aliev.

Extensively studied by scientists in the 1960s, the cave is considered to be the site of one of the most ancient proto-human habitations in Eurasia (a Neanderthal-style jaw bone - found in 1968 - is thought to be over 300,000 years old and thus one of the oldest proto-human remains found in this part of the world. Its discovery gave rise to the term Azykh Man). Despite of the few human remains discovered, evidence shows that the area was occupied by hominids over a period of nearly two million years.

The poor quality of the 1960's excavations led to uncertainty over the chronological position of the layers. In the Azykh cave 3rd layer (correlated with the Early Mousterian culture), the fossil fauna discovered consists of the following species: *Spaelacetus spelaeus*, *Ursus aff. arctos*, *Sus scrofa*, *Capreolus capreolus*, *Dama cf. mesopotamica*, *Megaloceros giganteus*, *Cervus elaphus*, *Capra aegagrus*, *Equus hydruntinus*, *Dicerorhinus merckii* (ALEV 1969).

Kazakhstan, § 2.5, pp. 26-28

According to TLEUBERDINA & NAZYMBETOVA (2010), at present "the *Elasmotherium sibiricum*, *Equus (Equus) mosbachensis*, *E. (Hemionus) hydruntinus*, *Dicerorhinus kirchbergensis* Koshkurgan horizon may be correlated with the end of the Lower Neopleistocene and the Middle Neopleistocene (around 500 kys BP)".

Several other *S. kirchbergensis* dental - as well as non-dental remains - have recently been discovered at the Koshkurgan village (14.5 Km north-east of Turkestan, Karatau Mountain, Chimkent region, South-Eastern Kazakhstan) (unpublished material, pers. communication). From this locality come other well-preserved twelve *S. kirchbergensis* isolated teeth (KHISAROVA 1963).

China, § 2.7, pp. 28-29

On the Anping locality (Liaoning, Liaoyang, North-eastern China), Xu (1986) apart, other *S. kirchbergensis* remains such as a maxilla (LA 7701-424), a mandibula (LA 7701-429), and a fragment of maxilla (LA 7701-521) - all preserving the teeth - were reported by ZHANG et al. (1980: Pl 2 - figs 1a, 1b, 1c, 1d).

From Yingkou near the Bohai Sea (Jinniu mountain, Liaoning province, North-eastern China) come few other *S. kirchbergensis* dental remains (JIN NIU SHAN EXCAVATION TEAM 1976: pp. 122-123; Pl I - figs 3, 4, 5).

A third lower premolar (5L167) comes from the Palaeolithic site of Lingjing (Xuchang, Henan Province, Central-Eastern China) (LI & DONG 2007; fig 4, n° 5) (Fig. 8, in this paper).

From the Migong Cave (Wushan County, Chongqing, Central China) - a late Late Pleistocene palaeoanthropological site (absolute age = 13150 ± 190 BP) - come




Fig. 8 - Third lower premolar of *S. kirchbergensis* from the Palaeolithic site of Lingjing (Xuchang, Henan Province, Central-Eastern China) (after LI & DONG 2007).
- Terzo premolare inferiore di *S. kirchbergensis* proveniente dal sito paleopolitico Lingjing (Xuchang, Provincia di Henan, Cina centro-orientale) (da LI & DONG 2007).




Fig. 9 - A maxillary fragment with three teeth ascribed to *S. kirchbergensis*, from the Tuozidong karst cave deposits (Jiangning District, Nanjing Municipality, Eastern China; after DONG et al. 2013).
- Un frammento di mascella con tre denti, riferito a *S. kirchbergensis*, proveniente dai depositi della Grotta di Tuozidong (Distretto di Jiangning, Municipalità di Nanjing, Cina orientale; da DONG et al. 2013).

cranial fragments, mandibles, teeth and some postcranial bones identified as *S. kirchbergensis*, representing the latest as well as the southernmost record of this species on the whole of Eurasia (CHEN et al. 2012). The related paper is rich in illustrations.

According to the authors (CHEN et al. 2012), "*S. kirchbergensis* migrated to the Yangtze River Valley during the Mid-Late Pleistocene probably because of the cold climate of the Ice Age. Based on *S. kirchbergensis* materials from the Migong Cave and other localities in the Yangtze River Valley, obvious distinctions are found among *S. kirchbergensis* from northern and southern China, and Europe. As a result, this species is divided into three types. Pleistocene rhinos from southern China, especially *Rhinoceros sinensis*, are also clarified".

From the Tuozidong karst cave deposits (Tuozijian, Humpback Cave, Humpback Hill, Jiangning District,




图 4 交界洞穴出土的奇蹄类和偶蹄类标本

梅氏犀 (*Stephanorhinus kirchbergensis*) 上颌骨残段 (97AJ:a61) 颚侧视 (A) 和嚼面视 (B), 右下颌骨 (97AJ:a62) 嚼面视 (C) 和颊侧视 (D), 左下颌骨 (97AJ:a62) 嚼面视 (E) 和舌侧视 (F); 北京斑鹿 (*Cervus (S.) nippon hortulorum*) 左下颌骨残段 (96AJ:17) 嚼面视 (G); 东北狍 (*Capreolus capreolus manchuricus*) 脱落角 (97AJ:19) 侧视 (H)

Fig. 4 Perissodactyl and artiodactyls from the Jiaojie Cave

Fig. 10- Mammals remains from the Jiaojie site (Heilongjiang province, North-eastern China): A-F indicate the *S. kirchbergensis* teeth (after YU & DONG 2011).

- Resti di mammiferi dal sito di Jiaojie (provincia di Heilongjiang, Cina nord-orientale): con A-F sono indicati i denti di *S. kirchbergensis* (da YU & DONG 2011).

Nanjing Municipality, Eastern China; GPS: 32° 03' 17.1" N, 119° 01' 57.6" E, 114 m asl) comes a maxillary fragment with three teeth ascribed to *S. kirchbergensis* (DONG et al. 2013: Fig. 9, in this paper). The great amount of the fossil mammal fauna collected together with the *S. kirchbergensis* remains were dated as Early Pleistocene (FANG et al. 2007).

From the Jiaojie site - discovered in 1996 - located in a karst cave (183 m asl) at Acheng (Haerbin municipality, Heilongjiang province, North-eastern China) come some *S. kirchbergensis* teeth (Fig. 10, in this paper). Up to now, this site represents the oldest palaeolithic site in the province as well as the northermost one in China (45° 21' 07.7" N - 127° 05' 16.8" E). The deposit is divided into 6 layers. The *S. kirchbergensis* teeth come from the 6th layer composed by reddish yellow clay with breccia. Uranium-series dating performed on the *S. kirchbergensis* teeth gave an age of 175 Ka BP (YU & DONG 2011).

An extraordinary well-preserved *S. kirchbergensis* skull has been discovered in the Heilongjiang province (North-eastern China) (unpublished material, pers. communication). No further data are presently available.

Just as also underlined by TONG & MOIGNE (2000), *Dicerorhinus* (= *Stephanorhinus*), *Coelodonta* and *Elasmotherium* are the rhinoceros species which mostly occurred in Northern China during Pleistocene. *Stephanorhinus kirchbergensis* (*D. kirchbergensis*) appears to be very common on this areal. On the contrary, *Rhinoceros* is actually confined to Southern China.

From the bone deposit of Kuhsiangtun (a site near Harbin, Northern Manchuria, discovered in 1926 and excavated in 1931-34) come *Rhinoceros* sp. remains (possibly *R. merckii* JÄGER) (LOUKASHKIN 1937: p. 331 and 334).

According to TONG et al. (2014), a *S. kirchbergensis* mandible (TNP00315) comes from Daheigou, while a second one (SGT2:1) comes from Shigou where a *S. kirchbergensis* left femur (SGT2:2) has been also found. "Mandibles and femur (fig 2-A, B, and C) (Fig. 11, in this paper) from Daheigou and Shigou were collected from Middle and Early Pleistocene strata, respectively. (omissis). The *S. kirchbergensis* fossils herein reported represent the definite earliest record with certain provenance of this species in China" (TONG et al. 2014: p. 338).



Fig. 11- *Stephanorhinus kirchbergensis* remains from the Nihewan Basin. A (TNP00315) mandible from Daheigou; B. (SGT2:1) mandible from Shigou; C. (SGT2:2) left femur from Shigou; A1, B1, buccal views. A2, B2, occlusal views; A3, B3, ventral views; C1, anterior view; C2, posterior view; C3, lateral view; C4, medial view (after TONG et al. 2014: fig. 2).

- Resti di *Stephanorhinus kirchbergensis* dal Bacino di Nihewan. A (TNP00315) mandibola da Daheigou; B. (SGT2:1) mandibola da Shigou; C. (SGT2:2) femore sinistro da Shigou; A1, B1, vista boccale. A2, B2, vista occlusale; A3, B3, vista ventrale; C1, vista anteriore; C2, vista posteriore; C3, vista laterale; C4, vista mediale (da TONG et al. 2014: fig. 2).

Lebanon and Israel, § 3.1, p. 29

The rhinoceros remains which “can unhesitatingly be referred to Merck’s rhinoceros, *Dicerorhinus merckii* (Jäger)” (= *S. kirchbergensis*) (HOOIJER 1961: p. 32) coming from Ksâr’ Akil (Lebanon) - “confined to levels XXVI A down to XXIX A” - consist of twenty-five upper teeth: a first and a third molars from the level XXVIII B, F4 are figured in Pl I - fig 3; a second deciduous molar from the level XXVII B, F5 is figured in Pl I - fig 4; a third premolar from the level XXVII B, F4 is figured in Pl I - fig 5; a third molar from the level XXVIII A, F5 is figured in Pl I - fig 6. Details and measurements of all the teeth are given in tabs 12-13.

The rhinoceros remains from Mugharet el-Emireh (BATE 1927: pp. 12-13) as well as those from Tabun-Mount Carmel (Israel) and the Skhul Cave (on the

slopes of Mount Carmel) (BATE 1937: p. 221 - figs 8j and 8m) were referred to *Rh. cf. hemitoechus* Falc., but - according to some other authors (cf. SCHROEDER 1903: p. 105; BERNSEN 1927: p. 25; SCHROEDER 1930: p. 7; ARAMBOURG 1958: p. 185) - they must be assigned to *R. merckii* (= *S. kirchbergensis*).

The rhinoceros from Oumm Qatafa near Jerusalem (Israel) was attributed by VAUFREY (1931: p. 254, 1951: p. 199) and later by WOLDSTEDT (1958) to “*Rh. Merckii*”.

Manuscript received on 10.I.2015, accepted on 22.I.2015

References

- AARIS-SØRENSEN, K. 1988. *Danmarks forhistoriske dyreverden*. Copenhagen: Gyldendals Bogklub, 252 pp.
- AARIS-SØRENSEN, K. 2009. Diversity and dynamics of the mammalian fauna in Denmark throughout the last glacial-interglacial cycle, 115-0 kyr BP. *Fossil and Strata* 57: 1-59.
- ALEKSEEV, M.N. 1970. An occurrence of Tiraspolian fauna at the Vilyuy River (Eastern Siberia). *Paleogeography, Paleoceanography, Paleoecology* 8: 209-14.
- ALEKSEEEVA, E.V. 1980. *Mlekopitayushchie Pleystozena Yugo-Vostoka Zapadnoy Sibiri (Hishchnye, Khobotnye, Kopytnye)* [Pleistocene Mammals from South-East of Western Siberia] [in Russian]. Moskva, Akademya Nauk SSSR, Dalnevostochny Nauchny Zentr Biologii (Pochv. Institut), Izd-vo “Nauka”: 187 pp.
- ALEV, S.D. 1969. *Fauna Paleoliticheskogo Mestonakhozhdennyia v Azikhe* [The Fauna of the Palaeolithic Site of Azikh] [in Russian]. PhD thesis. Baku: Izd-vo AN AzerbSSR (otdelenie Paleobiologii).
- ARAMBOURG, C. 1958. Les gros mammifères des couches tayaciennes. In *La Grotte de Fontéchevade, Part 3*, cur. H.V. VALLOIS, H. ALIMEN, C. ARAMBOURG & A. SCHREUDER, 185-225. Archives de l’Institut de Paléontologie Humaine, Mémoires 29.
- BATE, D.M.A. 1927. On the animal remains obtained from the Mugharet-El-Emireh in 1925. In *Researches in Prehistoric Galile*, cur. F. TURVILLE-PETRE, 9-13. London: British School of Archaeology in Jerusalem.
- BATE, D.M.A. 1937. Palaeontology - The fossil fauna of the Wady El-Mughara Caves. In *The Stone Age of Mount Carmel*, 137-240. Palaeontology 1, n. 2.
- BELYAEVA, E.I., & A.I. DAVID. 1975. Die Nashörner (Rhinocerotidae) aus dem Faunenkomplex von Tiraspol [with Russian & English abstracts]. *Quartärpaläontologie* 1: 157-75.
- BERKHOLST, B.E. 2011. *The large mammal fauna of the Pleistocene site Schöningen 13II - The levels Schö 13II-1, 13II-2 and 13II-3*. M. Sc. Thesis. Faculty of Archaeology, University of Leiden: 122 pp. [<http://www.rhinoresource-center.com/>].
- BERNSEN, J.J.A. 1927. *The Geology of the Tiglian Clay and its Fossil Remains of Rhinoceros*. Academisch Proefschrift [Ph.D. Thesis], Univ. van Amsterdam, C.N. Teulings’ Koninklijke Drukkerijen: 108 pp.
- BILLIA, E.M.E. 2008. Revision of the fossil material attributed to *Stephanorhinus kirchbergensis* (Jäger, 1839)

- (Mammalia, Rhinocerotidae) preserved in the museum collections of the Russian Federation. *Quaternary International*, 179, n. 1: 25-37 [http://dx.doi.org/10.1016/j.quaint.2007.09.034].
- BILLIA, E.M.E. 2011a. Occurrences of *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Rhinocerotidae) in Eurasia - An account. *Acta Palaeontologica Romaniae* 7: 17-40.
- BILLIA, E.M.E. 2011b. Siti paleontologici a "Rinoceronte di Merck", *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Perissodactyla), in Istria, Quarnero e Dalmazia [Paleontoloska Najdisca Merckovega Nosoroga, *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Perissodactyla) v Istri in Dalmaciji - Paleontoloska nalazista Merckovog Nosoroga, *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Perissodactyla), u Istri i Dalmaciji] [with Slovenian & Croatian summ.]. *Atti Centro Ricerche Storiche, Rovigno / Centar za Povijesna Istrazivanja, Rovinj / Središće za Zgodovinska Raziskovanja Rovinj* 41: 9-31.
- BILLIA, E.M.E. 2014. *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Rhinocerotidae) from European Russia - A new, detailed inventory of sites and referred material. *Central European Geology* 57, n. 2: 165-95 [doi: 10.1556/CEuGeol.57.2014.2.4].
- BREDA, M., & M. MARCHETTI. 2003. I mammiferi fossili - *Stephanorhinus* cfr *etruscus* (Falconer, 1859), S. cfr *hundsheimensis* (Toula, 1902), S. *kirchbergensis* (Jäger, 1834) (supervisione di B. Sala). In *Gli Antichi Bacini Lacustri e i Fossili di Leffe, Ranica e Pianico-Sellere, Prealpi Lombarde*, cur. C. RAVAZZI, 44-8. Quaderni Geodinamica Alpina e Quaternaria CNR - Quaderni della Comunità Montana Valle Seriana 2.
- BREDA, M., & M. MARCHETTI. 2007. Pleistocene mammals faunas from the Leffe Basin (Bergamo, Northern Italy) - Revision and new data. *Courier Forschungs-Institut Senckenberg* 259: 61-77.
- CEPEK, A.G. 1986. Schichtenfolge und Position des "Rixdorfer Horizontes". In *Kurzreferate und Exkursionsführer „25 Jahre AK Quartärgeologie“*, 19-21. Berlin: Gesellschaft für Geologische Wissenschaften der DDR.
- CHEN, S.-K., W.-B. HUANG, J. PEI, C.-D. HE, L. QIN, G.-B. WEI & J. LENG. 2012. The latest Pleistocene *Stephanorhinus kirchbergensis* from the Three Gorges Area and re-evaluation of Pleistocene rhinos in Southern China. *Acta Anthropologica Sinica*, 31, n. 4: 381-94.
- CHEPALYGA, A.L. 1967. *Antropogenovye Presnovodnye Mollyuski Yuga Russkoy Ravniny i ikh Stratigraficheskoe Znechenie* [Anthropogene Freshwater Mollusks in the South of the Russian Plain and their Stratigraphic Importance] [in Russian]. Moskva, Trudy Akad. Nauk SSSR 166: 222 pp.
- CHERSKY, I.D. 1878. Predvaritel'noe soobshchenie o dostavlennoj iz Verkhoyanskogo Okruga golovy nosoroga (*Rhinoceros antiquitatis* s. *tichorhinus*) s" [in Russian]. *Izvestya Vostochno-Sibirskago otdela Imperatorskago Russkago geograficheskago ob-va* 9, n. 5-6: 165.
- CHERSKY, I.D. 1879. Opisanie golovy sibirskago nosoroga (*Rhinoceros antiquitatis* s. *tichorhinus*), naydennoy v" 1877 godu v" Verkhoyanskem" okrug" s" sokhranivshimisya pri ney myagkimi chastyami [Description of a Siberian rhinoceros skull found in 1877 in the neighbourhood of Verkhoyansk ...] [in Russian]. *Izvestya Vostochno-Sibir-* skago otdela Imperatorskago Russkago geograficheskago ob-va 110, n. 1-2: 36-59.
- CZARNECKI, A. 1983. Zur Entwicklung des Menschen in Südwestdeutschland. In *Urgeschichte in Baden Württemberg*, cur. H. MÜLLER-BECK, 217-40. Stuttgart: Thesis.
- DEGERBØL, M.A. 1952. Recent discoveries of remains of mammals from glacial deposits in Denmark: *Dicerorhinus kirchbergensis* (Jäger), new to the fauna of Denmark; *Megaloceros giganteus* (Blumenbach) and *Bison priscus* (H. v. Meyer). *Det Kongelige Danske Videnskabernes Selskab-Biologiske skrifter* 6, n. 8: 1-26.
- DIEDRICH, C.G., & K. ZAK. 2006. Prey deposits and den sites of the Upper Pleistocene hyaena *Crocuta crocuta spelaea* (Goldfuss, 1823) in horizontal and vertical caves of the Bohemian Karst (Czech Republic). *Bulletin of Geosciences* 81, n. 4: 237-76.
- DIETRICH, W.O. 1932. Über den Rixdorfer Horizont im Berliner Diluvium. *Zeitschrift der Deutschen Geologischen Gesellschaft* 84, n. 4: 193-221.
- DONG, W., J. LIU & Y.S. FANG. 2013. The large mammals from Tuozidong (eastern China) and the Early Pleistocene environmental availability for early human settlements. *Quaternary International* 295: 73-82 [doi:10.1016/j.quaint.2012.01.014].
- DORONICHEV, V. 2008. The Lower Paleolithic in Eastern Europe and the Caucasus - A Reappraisal of the Data and New Approaches. *PaleoAnthropology* 2008: 107-157.
- DUBROVO, I.A. 1957. Ob ostatkakh *Parelephas wüsti* (Pawl.) i *Rhinoceros mercki* Jaeger iz Yakutii [On remains of *Parelephas wüsti* (Pawl.) and *Rhinoceros mercki* Jaeger from Yakutya] [in Russian]. *Byulleten' Komissii po Izuchenju Chetvertichnogo Perioda* 21: 97-104.
- ĐURIŠOVÁ, A. 1994. *Dicerorhinus kirchbergensis* (Mammalia, Rhinocerotidae) z fluvialnych náplavov Váhu v Šal'i (Slovenská republika) [*Dicerorhinus kirchbergensis* (Mammalia, Rhinocerotidae) from the fluvial deposits Vahu at Šal'i (Slovakian Republic)]. *Zborník Slovenského národného Muzea (Prírodné Vedy)* 40: 7-13.
- FABIANI, R. 1919. I Mammiferi Quaternari della Regione Veneta. *Memorie dell'Istituto di Geologia della Regia Università di Padova* 5: 1-174.
- FANG, Y.-S., W. DONG et al., cur. 2007. *The Early Pleistocene Mammalian Fauna at Tuozi Cave, Nanjing, China*. Beijing, Science Press (in Chinese, English summ.).
- FRIEDEL, E. 1880. Ein Leichnam von *Rhinoceros Merckii* aus Sibirien. *Der Zoologische Garten - Zeitschrift für Beobachtung, Pflege und Zucht der Thiere* 21, n. 12: 353-9.
- GORJANOVIĆ-KRAMBERGER, D. 1904. Der paläolithische Mensch und seine Zeitgenossen aus der Diluvium von Krapina in Kroatien. *Mitteilungen der Anthropologischen Gesellschaft* 31: 189-216.
- GORJANOVIĆ-KRAMBERGER, D. 1906. *Der Diluviale Mensch von Krapina in Kroatien*. Wiesbaden, Kreidels: 277 pp.
- GORJANOVIĆ-KRAMBERGER, D. 1913a. Zivot i kultura diluvijalnoga covjeka iz Krapine u Hrvatskoj. *Djela Jugoslavenske Akademije (JAZU)* 23: 1-54.
- GORJANOVIĆ-KRAMBERGER, D. 1913b. Fossilni rinocerotidi Hrvatske i Slavonije, s osobitim obzirom na *Rhinoceros Mercki* iz Krapine (De rhinocerotidibus fossilibus Croatie et Slavoniae, praecipua ratione habita *Rhinocerotis Mercki* var. *Krapinensis* mihi) [in Croatian and Latin]. *Djela Jugoslavenske Akademije (JAZU)* 22, n. 8: 1-70.

- GROMOVA, V.I. 1935. Ob ostatkakh nosoroga Merka (*Rhinoceros merckii* Jaeg.) s Nizhney Volgi [Ueber die Reste des Merckschen Nashorn (*Rhinoceros merckii* Jaeg.) von der unteren Wolga] [in Russian, German abstr.]. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR* 4: 91-131.
- GÜRICH, G. 1905. Der Scheckenmergel von Ingaramsdorf und andere Quartärfunde in Schlesien. *Jahrbuch der Königlich Preussischen Geologischen Landesanstalt und Bergakademie* 26: 43-57.
- HEINRICH, W.-D. 2002. Der Rixdorfer Horizont - Ein Fundstättentyp eiszeitlicher Säugetiere in Berlin und Brandenburg. *Humboldt-Spektrum* 2-3: 70-5.
- HERMANN, R. 1911. Über *Rhinoceros Merckii* Jäger im Diluvium Westpreußens und seine Beziehungen zur norddeutschen Diluvialfauna. *Zeitschrift der Deutschen Geologischen Gesellschaft (Abhandlungen und Monatsberichte)* 63, n. 1: 13-35.
- HERMSDORF, N. 2000. Die Sandgrube Niederlehme - Ein klassischer Aufschluss des Rixdorfer Horizontes. *Brandenburger Geowissenschaftliche Beiträge* 7: 173-81.
- HERMSDORF, N. 2002. Neue Daten zur Altersstellung des Rixdorfer Horizontes von Niederlehme. *Brandenburger Geowissenschaftliche Beiträge* 1-2: 44.
- HOOIJER, D.A. 1961. The fossil vertebrates of Ksâr' Akil, a palaeolithic rock shelter in the Lebanon. *Zoologische Verhandelingen* 49: 3-68.
- JELINEK, J. 1969. Neanderthal man and *Homo sapiens* in Central and Eastern Europe. *Current Anthropology* 10: 475-503.
- JIN NIU SHAN EXCAVATION TEAM, 1976. [Significance of Quaternary mammalian fauna found in Yingkou, Liaoning]. *Vertebrata PalAsiatica* 14, n. 2: 120-7.
- KAFKA, J. 1913. Rezente und fossile Huftiere Böhmens (Ungulata). *Archiv für die Naturwissenschaftliche Landesforschung von Böhmen* 14, n. 5: 1-86.
- KHISAROVA, G.D. 1963. Iskopaemye kosti mlekopitayushchikh iz Koshkurganskogo grifona, Yuzhny Kazakhstan. In *Materialy po Istorii Fauny i Flory Kazakhstana*, 42-65 [Mammal osteological fossil remains from the Koshkurgan locality, Southern Kazakhstan (in Material on the History of the Fauna and Flora of Kazakhstan) [in Russian]. Akademya Nauk Kazakhstanskoy SSR (Institut Zoologii), Izd-vo "Nauka" KazSSR 4.
- KRUMBECK, L. 1917. Beiträge zur Geologie von Nordbayern - I. Über einen Zahn von *Rhinoceros* aus dem Diluvium von Spardorf bei Erlangen. *Sitzungsberichte der Physikalisch-Medicinische Sozietät zu Erlangen* 48: 124-30.
- LANSER, K-P. 1997. Der Schädel eines dicerorhininen Nashorns aus der Dechenhöhle bei Iserlohn-Letmathe. *Geologie und Paläontologie in Westfalen* 47: 53-78.
- LEONARDI, P. 1945-47. Resti fossili di rinoceronti del Museo di Storia Naturale di Trieste. *Atti del Museo Civico di Storia Naturale di Trieste* 16, n. 12: 145-60.
- LEONARDI, P. 1947. Resti fossili inediti di rinoceronti conservati nelle collezioni dell'Istituto Geologico dell'Università di Padova. *Memorie dell'Istituto di Geologia dell'Università di Padova* 15: 1-30.
- LI, Z.Y., & W. DONG. 2007. Mammalian Fauna from the Lingjing Paleolithic site in Xuchang, Henan Province [in Chinese, English abstr.]. *Acta Anthropologica Sinica* 26 (4): 345-60.
- LOBACHEV, Yu.V., A.Yu. LOBACHEV & E.M.E. BILLIA. 2014. Morfologicheskie i biomechanicheskie osobennosti zhevatel'nogo aparata *Coelodonta antiquitatis* (Blumenbach, 1799) i *Stephanorhinus kirchbergensis* (Jäger, 1839) [Morphological and biomechanical features of the masticatory apparatus in *Coelodonta antiquitatis* (Blumenbach, 1799) and *Stephanorhinus kirchbergensis* (Jäger, 1839)] [in Russian]. *Materialy LX sessii Paleontol. Obstva "Diversifikazyi i Etapnost' Evolyuzii Organicheskogo Mira v Svetle Paleontologicheskoy Letopisi"* 60: 168-70.
- LONA, F. 1963a. Floristic and glaciologic sequence (from Donau to Mindel) in a complete diagram of the Leffe deposit. *Berichte Geobotanischen Institutes Eidg. Technischen Hochschule* 34: 64-6.
- LONA, F. 1963b. Comparative diagrams of some Pliocene-Pleistocene marine sediments in the Po-Valley and the continental deposits of Leffe (Bergamo). *Berichte Geobotanischen Institutes Eidg. Technischen Hochschule* 34: 66-7.
- LOUKASHKIN, A.S. 1937. Some observations on the remains of the Pleistocene fauna and the Palaeolithic age in northern Manchuria. In *Early Man International Symposium*, cur. G.G. MACCURDY, 327-340. Philadelphia: Academy of Natural Sciences, J.B. Lippincott Co.
- MALEZ, M. 1958. Neki noviji rezultati paleontoloskog istrazivanja Pećine Veternice [Einige neue Resultate der paläontologischen Erforschung der Höhle Veterna] [in Croatian & German]. *Jugoslovenska Akademija Znanosti i Umjetnosti (Odjel za Prirodne Nauke)*, 5-18 (in Croatian) and 19-24 (in German).
- MALEZ, M. 1961. Nalaz dvaju pleistocenskih sisavaca kod Zagreba i pregled okolnih nalazista [The find of two Pleistocene mammals near Zagreb and an overview on adjacent find localities] [in Croatian, German summary]. *Geoloski vjesnik* 14: 63-88.
- MALEZ, M. 1963a. Pećina Veterna u Medvednici - I. Opci speleoloski pregled - II. Stratigrafija kvarternih talozina. *Acta geologica* 5: 175-229.
- MALEZ, M. 1963b. Qvartarna Fauna Pećine Veternice u Medvednici [Die Quartäre Fauna der Höhle Veterna (Medvednica, Kroatien)] [in Croatian, German summary]. *Palaeontologia Jugoslavica* 5: 108-12+178-9.
- MALEZ, M. 1978a. Novija istrazivanja paleolitika u Hrvatskom Zagorju. In *Arheoloska istrazivanja u Sjeverozapadnoj Hrvatskoj*, cur. Z. RAPANIC, 6-69. Hrvatsko arheolosko drustvo 2.
- MALEZ, M. 1978b. Populacije neandertalaca i neandertalcima slicnih ljudi u Hrvatskoj. In *Krapinski Pracovjek i Evolucija Hominida*, cur. M. MALEZ, 331-71. Zagreb: Jugoslavenska Akademija Znanosti i Umjetnosti.
- MALEZ, M. 1978c. Fossile Menschen aus Nordwestkroatien und ihre quartärgeologische, paläontologische und paläolithische Grundlage. *Collegium Antropologicum* 2: 29-41.
- MALEZ, M. 1986. Die quartären Vertebraten-Faunen in der SFR Jugoslawien [English & Russian summ.]. *Quartärpaläontologie* 6: 101-17.
- MALEZ, M., F.H. SMITH, J. RADOVČIĆ & D. RUKAVINA, 1980. Upper Pleistocene hominids from Vindija, Croatia, Yugoslavia. *Current Anthropology* 21: 365-7.
- MIRACLE, P., & D. BRAJKOVIĆ. 1992. Revision of the Ungulate Fauna and Upper Pleistocene Stratigraphy of Vaternica Cave (Zagreb, Croatia). *Geologia Croatica* 45: 1-14.

- MOL, D., K. POST & H. v. d. PLICHT. 2012. Fossielen van bosneushoorn (*Stephanorhinus kirchbergensis*) en bosolifant (*Elephas antiquus*) uit het Eurogeulgebied. *Cranium* 29: 20-5.
- MOSTEKY, V. 1966. Rhinocerotidae aus der Höhle "Chlupáčova Sluj" bei Koněprusy. *Sbornik Narodniho Musea v Praze - Acta Musei Nationalis Pragae* 22, n. 4: 143-61.
- NEHRING, A. 1878. Die quaternären Faunen von Thiede und Westeregeln. *Archiv für Anthropologie* 10: 359-98, 11: 1-24.
- NEHRING, A. 1880. Übersicht über vierhundzwanzig mittel-europäische Quartär-Faunen. *Zeitschrift der Deutschen Geologischen Gesellschaft* 32: 473.
- PENKMAN, K., M. COLLINS, D. KEEN & R. PREECE. 2008. *British Aggregates: an Improved Chronology of Using Amino Acid Racemisation and Degradation of Intracrystalline Amino Acids*. English Heritage Research Department Report Series 6/2008.
- PERSICO, D., E.M.E. BILLIA, S. RAVARA & B. SALA. 2014. The skull of *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Rhinocerotidae) from Spinadesco (Cremona, Lombardia, Northern Italy): morphological analyses and taxonomical remarks - An opportunity for revising the three other skulls from the Po Valley. *Quaternary Science Reviews* 109: 28-37 [doi.org/10.1016/j.quascirev.2014.11.022].
- POHLIG, H. 1887. Über *Elephas trogontherii* und *Rhinoceros Merckii* von Rixdorf bei Berlin. *Zeitschrift der Deutschen Geologischen Gesellschaft* 39: 798-807.
- RAKOVEC, I. 1942. Nosorog s Kamnitnika pri Skofji Loki [On the rhinoceros remains from Kamnitnik hill near Skofja Loka] [in Slovenian]. *Mat.-Prirodoslov. Razprave [Math.-Natural History Dissertation]*. Slovenka Akademija Znanosti i Umetnosti 2: 241-62.
- RAKOVEC, I. 1958. Les Mammiferes pléistocenes de l'Abrib "Crvena Stijena" près de Petrovici (Monténégro). *Glasnik Zemaljskog Muzeja* n.s., 13: 65-75.
- RAKOVEC, I. 1959. Kvartarna sesalka favna iz Betalovega spodmola pri Postojni [Quaternary mammalian fauna from the Betalov Spodmol Cave near Postojna]. *Razprave* 5, Kl. IV: 287-348.
- RAVAZZI, C., R. PINI & M. BREDA. 2009. Reconstructing the palaeoenvironments of the Early Pleistocene mammal faunas from the pollen preserved on fossil bones. *Quaternary Science Reviews* 28: 1-15.
- REEKER, H. 1906-07. *Rhinoceros merckii* Jäger in Westfalen. *Fünfunddressigster Jahresbericht des Westfälischen Provinzial-Vereins für Wissenschaft und Kunst*: 31-2.
- SATAEV, R.M. 2008. Ostatki medvedya Ursus (Spelaeartos) sp. i nosoroga Stephanorhinus sp. iz peshchery Verkhnyaya (YUzhny Ural) [The remains of bear Ursus (Spelaeartos) sp. and rhinoceros Stephanorhinus sp. from the cave "Verkhnyaya" ("Upper") (Southern Ural)]. In *Fauna i Flory Severnoy Evrazii v Pozdнем Кайноzoе - Faunae and Flora of Northern Eurasia in the Late Cenozoic*, cur. P.A. KOSINTSEV, 137-46. Ekaterinburg/Chelyabinsk: Rossijskaya Akademya Nauk, Ural'skoe otdelenie, Institut Ekologii Rastenii i Zhivotnikh.
- SCHMIDT, Z. 1969. Nálezy chladnomilných nosorožcov na Slovensku. *Svet vedy, Slovenska Akademia Vied* 16, n. 9: 536-40.
- SCHRENCK, L.v. 1880. Der erste Fund einer Leiche von *Rhinoceros merckii*. *Zapiski Imperatorskoy Akademii Nauk* [Mémoires de l'Académie Impériale des Sciences de St.-Pétersbourg] s. 7, 28, n. 7: 1-55.
- SCHREVE, D.C. 1997. *Mammalian Biostratigraphy of the Later Middle Pleistocene in Britain*. Unpublished PhD thesis, University College of London.
- SCHROEDER, H. 1903. Die Wirbeltier-Fauna des Mosbacher Sandes - 1. Gattung *Rhinoceros*. *Abhandlungen der Kaiserlich-Königlichen Preussischen Geologischen Landesanstalt* N.F., 18: 1-143.
- SCHROEDER, H. 1905. *Rhinoceros merckii* Jäger von Heggen im Sauerlande. *Jahrbuch der Königlich Preussischen Geologischen Landesanstalt Bergakademie* 26, n. 2: 212-39.
- SCHROEDER, H. 1930. Über *Rhinoceros merckii* und seine nord- und mitteldeutschen Fundstellen. *Abhandlungen der Königlich Preussischen Geologischen Landesanstalt* n.f., 124: 1-114.
- SHPANSKY, A.V., & E.M.E. BILLIA. 2012. Records of *Stephanorhinus kirchbergensis* (Jäger, 1839) (Mammalia, Rhinocerotidae) from the Ob' River at Krasny Yar (Tomsk region, southeast of Western Siberia) [with Russian abstract]. *Russky Teriologichesky Zhurnal - Russian Journal of Theriology* 11, n. 1: 47-55.
- SINZOW, J. 1900. Geologische und paläontologische Beobachtungen in Südrussland. *Zapiski Imperatorskago Novorossiyskago Universiteta* 79: 347-412.
- SLÓSARSKI, A. 1884. Zwierzeta zaginione (dyluwialne) - Kilka słów o nosorozcach dyluwialnych - *Rhinoceros tichorhinus* Fischer i *Rhinoceros Merckii* Jäger [Extinct animals (diluvial) - Some words on diluvial rhinoceroses - *Rhinoceros tichorhinus* Fischer and *Rhinoceros Merckii* Jäger] [in Polish]. *Pamietnik Fizyjograficzny* 4, n. 4: 357-74.
- SMITH, F.H. 1976a. A fossil hominid frontal from Velika Pećina (Croatia) and a consideration of Upper Pleistocene hominids from Yugoslavia. *American Journal of Physical Anthropology* 44: 127-34.
- SMITH, F.H. 1976b. *The Neandertal Remains from Krapina - A Descriptive and Comparative Study*. University of Tennessee Department of Anthropology, Report of Investigations 15.
- STAESCHE, K. 1941. Nashörner der Gattung *Dicerorhinus* aus Diluvium dem Würtemberg. *Abhandlungen der Reichsstelle für Bodenforschung* 200: 1-148.
- STEHLIN, H.G. 1930. Die Säugetierfauna von Leffe (prov. Bergamo). *Eclogae Geologicae Helvetiae*, 23 (2): 648-81.
- STEPANOV, V.V. 1961. Chetvertichnye otlozheniya v dolinakh limanov Odesskoy oblasti [Quaternary deposits in dolines of estuaries in the Odessa oblast]. In *Materialy Vsesoyuznogo Soveshchaniya po Izuchenju Chertvertichnogo Perioda. Tom 2*. Moskva.
- TARAMELLI, T. 1898. Del deposito lignitico di Leffe in provincia di Bergamo. *Atti della Società Geologica Italiana* 17: 202-18.
- TLEUBERDINA, P.A., & G. NAZYMBETOVA. 2010. Distribution of *Elasmotherium* in Kazakhstan. In *Quaternary Stratigraphy and Paleontology of the Southern Russia - Connections between Europe, Africa and Asia*, cur. V.V. TITOV & A.S. TESAKOV, 171-3. Rostov-on-Don: Abstracts of the International INQUA-SEQS Conference, Rostov-on-Don, June 2010, Southern Scientific Centre RAS [http://seqs2010-rostov.narod.ru/absractsrostovpdf, http://www.rhinoresourcecenter.com/ref_files/1303341272.pdf].

- TONG, H., & A.-M. MOIGNE. 2000. Quaternary rhinoceros of China. *Acta Anthropologica Sinica* 19 (suppl.): 257-63.
- TONG, H., F. WANG, M. ZHENG & X. CHEN. 2014. New fossils of *Stephanorhinus kirchbergensis* and *Elasmotherium peii* from the Nihewan Basin. *Acta Anthropologica Sinica* 33 (3): 369-88.
- TOULA, F. 1902. Das Nashorn von Hundsheim, *Rhinoceros (Ceratotherinus Osborn) hundsheimensis* nov. form. mit Ausführungen über die Verhältnisse von elf Schädeln von *Rhinoceros (Ceratotherinus) sumatrensis*. *Abhandlungen der Kaiserlich-Königlichen Geologischen Reichsanstalt* 19, n. 1: 1-92.
- TOULA, F. 1907. *Rhinoceros merckii* Jäger in Österreich. *Jahrbuch der Kaiserlich-Königlichen Geologischen Reichsanstalt* 57, n. 3: 445-54.
- TYRÁČEK, J., O. FEJFAR, J. FRIDRICH, J. KOVANDA, L. SMOLÍKOVÁ & J. SÝKOROVÁ. 2001. Račiněves - A new Middle Pleistocene interglacial in the Czech Republic. *Bulletin of the Czech Geological Survey* 76: 127-39.
- TYRÁČEK, J., R. WESTAWAY & D.R. BRIDGLAND. 2004. River terraces of the Vltava and Labe (Elbe) system, Czech Republic, and their implications for the uplift history of the Bohemian Massif. *Proceedings of the Geologists' Association* 115: 101-24.
- VALOCH, K. 1968. Evolution of the Paleolithic in Central and Eastern Europe. *Current Anthropology* 9: 351-390.
- VAUFREY, R. 1931. L'Acheuléen supérieur de la Grotte d'Oumm-Qatafa (Palestine) - IV. Paléontologie. *L'anthropologie* 41: 253-63.
- VAUFREY, R. 1951. Mammifères. In *Le Paléolithique et le Mesolithique du Desert de Judee*, cur. R. NEUVILLE et al., 198-217. Archives de l'Institute Paléontologie Humaine, Memoires 24.
- VENZO, S. 1956. Stadi della glaciazione del "Donau" sotto al Günz e al Mindel nella serie lacustre di Leffe (Bergamo) - Limite tra Pliocene e Pleistocene. *Atti del IV Congresso Internazionale del Quaternario* 1: 65-85.
- VIALLI, G. 1956. Sul rinoceronte e l'elefante dei livelli superiori della serie lacustre di Leffe (Bergamo). *Memorie della Società Italiana di Scienze Naturali (Museo Civico di Storia Naturale di Milano)* 12, n. 1: 1-71.
- VUKOVIĆ, S. 1953. Pecina Vindija kao preistorijska stanica. *Speleolog-Časopis Speleološke Sekcije P.D. Željezničar* 1, n. 1: 15-23.
- VUKOVIĆ, S. 1954. Istrazivanje pretprecinskog terena Vindije. *Speleolog-Časopis Speleološke Sekcije P.D. Željezničar* 2, n. 1: 23-8.
- WINGS, O., D. SCHWARZ-WINGS. & W.-D. HEINRICH. 2009. The Rixdorf Horizon ("Rixdorfer Horizont") of the Niederlehme sand pits near Königs Wusterhausen. *Abstracts volume of the 7th Annual EAVP Meeting 2009*: 71-4.
- WOLDRICH, J.N. 1882. Beiträge zur Fauna der Breccien und anderer Diluvial-gebilde Oesterreichs mit besonderer Berücksichtigung des Pferdes. *Jahrbuch der Kaiserlich-Königlichen Geologischen Reichsanstalt* 32 n. 4: 435-70.
- WOLDSTEDT, P. 1958. *Das Eiszeitalter*. Stuttgart, 2: 438 pp.
- WOLPOFF, M.H. 1979. The Krapina dental remains. *American Journal of Physical Anthropology* 50: 67-113.
- WOLPOFF, M.H. 1980. *Paleoanthropology*. New York: Knopf, 878 pp.
- WOLPOFF, M.H., F.H. SMITH, M. MALEZ, J. RADOVCIĆ & D. RUKAVINA. 1981. Upper Pleistocene human remains from Vindija Cave, Croatia, Yugoslavia. *American Journal of Physical Anthropology* 54: 499-545.
- WÜST, E. 1901. Untersuchungen über das Pliozän und das älteste Pleistozän Thüringens, nördlich vom Thüringer Wald und westlich von der Saale. *Abhandlungen der Naturforschenden Gesellschaft zu Halle* 23: 265-81.
- XU, X.-F. 1986. [Dicerorhinus kirchbergensis (Jäger, 1839) from the Late Middle Pleistocene mammalian fauna of Anping (Liaoning)] [in Chinese, English summary]. *Vertebrata PalAsiatica* 24, n. 3: 229-41.
- YU, H., & W. DONG. 2011. [Pleistocene mammalian fauna from the Jiaojie cave at Acheng, Heilongjiang, Province] [in Chinese, English abstr]. *Quaternary Sciences* 31 (4): 675-88 [doi:10.3969/j.iss.1001-7410.2011.04.011].
- ZHANG, Z.-H., B. ZOU & L. ZHANG. 1980. [The discovery of fossil mammals at Anping, Liaoyang] [in Chinese, English summary]. *Vertebrata PalAsiatica* 18, n. 2: 154-62.

Authors' addresses - Indirizzi degli Autori:

- Emmanuel M.E. BILLIA
Via Bacchiglione 3, I-00199 ROMA
e-mail: e.billia@yandex.ru, eme.billia@alice.it
- Julia ZERVANOVÁ
Drobného 16/9, SK-84101 BRATISLAVA
e-mail: lia20lia@gmail.com